

Recognizable and Rational Formal Power Series with General Discounting

Manfred Droste

Department of Mathematics and Computer Sciences

University of Leipzig

D-04109 Leipzig, Germany

E-mail: `droste@informatik.uni-leipzig.de`

Heiko Vogler

Department of Computer Science

Dresden University of Technology

D-01062 Dresden, Germany

E-mail: `vogler@tcs.inf.tu-dresden.de`

We investigate the behaviours of weighted automata with general discounting; these are the skew (formal power) series. Recently, it was shown that the recognizable skew series are precisely the rational skew series [DK03]; this generalizes a classical result of Kleene-Schützenberger [Kle56, Sch61]. Here, we show that a very special case of Schützenberger’s result suffices to deduce both its general version and the skew generalization.

References

- [DK03] M. Droste and D. Kuske. Skew and infinitary formal power series. In *ICALP 2003*, volume 2719 of *LNCS*, pages 426–438. Springer-Verlag, 2003.
- [Kle56] S. E. Kleene. Representation of events in nerve nets and finite automata. In C.E. Shannon and J. McCarthy, editors, *Automata Studies*, pages 3–42. Princeton University Press, Princeton, N.J., 1956.
- [Sch61] M.P. Schützenberger. On the definition of a family of automata. *Inf. and Control*, 4:245–270, 1961.

